



## **MAFAS LTD**

Lot 167, Bay View, Concord Road, La Romain. Trinidad W.I.  
Tel: 868-354-9666; Tel/Fax: 868-697-1375. Email: marcusmycoo@yahoo.com  
Visit us online: www.mafascaribbean.com

### **CABBAGE, CAULIFLOWER, BROCCOLI** **CRUCIFER PRODUCTION GUIDE**

Day1: Same day of transplanting apply as a drench at the base of the plant at a cool time:

**DRENCH** only: Do not apply over the plant.

**X-tra Power**(30 ml/gal) + **Harvest More 10:55:10**(2tsp/gal) + **CalGuard**(10ml/gal) + **Nitro-Plus 23**(15ml/gal) + Insecticide

**FOLIAR** apply:

5-7 DAT: **Nitro Plus 23**(30ml/gal) + **Vine Mix**(30ml/gal) + **Harvest More 10:55:10**(2tsp/gal) + **Top Cop**(15ml/gal)

10-14 DAT: **Nitro Plus 23**(30ml/gal) + **Harvest More 13:0:44**(2tsp/gal) + **Golden Pest Spray Oil**(30ml/gal)

15-21 DAT: **Folizyme**(30ml/gal) + **Harvest More 13:0:44**(2tsp/gal) + **Top Cop**(15ml/gal) + Insecticide

20-28 DAT: **Sett**(30ml/gal) + **Vine Mix**(30ml/gal) + **X-cyte**(5ml/gal) + **Harvest More 20:20:20**(2tsp/gal)

25-35 DAT: **Folizyme**(30 ml/gall) + **Harvest More 5:5:45**(2tsp/gal) + **Top Cop**(15ml/gal) + Insecticide

30-42 DAT: **Sett**(30ml/gal) + **Vine Mix**(30ml/gal) + **X-cyte**(5ml/gal) + **Harvest More 5:5:45**(2tsp/gal)

35-49 DAT: **Folizyme**(30ml/gal) + **Harvest More 5:5:45**(2tsp/gal) + **Top Cop**(15ml/gal)

40-56 DAT: **Sett**(30ml/gal) + **Harvest More 5:5:45**(2tsp/gal) + **Golden Pest Spray Oil**(30ml/gal)

**Soil applied Fertilizers-** At farmers discretion but consider reducing 25-30% rates and intervals; Starter fertilizer should be 12:24:12 and 2 weeks before heading at 14-21 days applications of 12:12:17 + 2 or 12:6:35 +2 or a similar "bearing salt". Soil applied Fertilizers may need to be adjusted as the products increase efficiency of uptake. Review fertilizer application intervals and quantity.

- Farmers are encouraged to contact **MAFAS LTD** personnel for further information on effective disease and pest control on crop production
- **Note well:** **Stimulate** may be applied 14-21 days after it is first used at 17-20 day.
- The soil surface below the leaves should also be sprayed with a soil acting contact fungicide early in the crop especially where manure is used.
- **Top Cop** may be applied at 30ml/gal for control of a wide range of leaf spots. Do a compatibility test. Can be mixed with **Sett**, **Nitro-plus 23**, **Stimulate** and **X-cyte**.
- Ensure good coverage when using **Golden Pest Spray oil**. **GPS** is approved for use in **ORGANIC CROPPING**
- Never mix **X-cyte** with **Stimulate**
- **Stimulate** is applied at knap sack rate of **5ml/4 gall**
- Mist blower rates are twice the knapsack rate.

#### **Technical Staff:**

**Marcus Mycoo: 360-1986; Perry Jones: 354-9668; Rayard Khan: 354-9665**