



MAFAS LTD

Lot 167, Bay View, Concord Road, La Romain. Trinidad W.I.
Tel: 868-354-9666; Tel/Fax: 868-697-1375. Email: marcusmycoo@yahoo.com
Visit us online: www.mafascaribbean.com

PROGRAMME FOR HOT PEPPER

Always use **High Quality, 3 week old healthy seedlings. Apply Rezist at 30 ml/gall to seedling week before transplanting to develop tolerance to viruses and other diseases.**

At Transplanting: Soil DRENCH at the base of the plant only for Damping Off etc: Do not apply over the leaves.

Diligent 81 CT (11/2Tbsp/gal) + Aldabon (15ml/gall) + Binder (5ml/gal) + X-Tra Power (30ml/gal) + Bioforge (5 ml/gall) + Harvest More 10:55:10 (2 tsp/gal)

5-7 days later: Drench the plants at the base of plants. Not over the leaves

Rotaprid (10ml/gal) + X-Tra Power (30ml/gal) + Phytoguard Cal (5 ml/gall) + Charge (30m/gall) + Harvest More 10:55:10 (1Tbsp/gal)

FROM 14 DAYS AFTER TRANSPLANTING and every 14-21 days thereafter: Fertigation Programme

***Root Feed SP 1kg/barrel + Charge 1L/barrel + 1 kg/barrel 5-10-27-10 Urea Mate-Harvest More**

FOLIAR and Drench apply: In Stoller pH+ treated water

7-10 days Later: **Diligent 81 CT (1 TBSp/gal) + Obulus (5.0 ml/gal) + Nitro-Plus (20 ml/gal) + Rezist (20ml/gal) + 10:55:10 HM (1Tbsp/gal)**

7-10 days: **Rodazim (5ml/gal) + Obulus (5 ml/gal) + Nitro-Plus (20 ml/gal) + Vine Mix (20 ml/gal) + Bio-Forge (10ml/gal) + 10:55:10 HM (1Tbsp/gal)**

7-10 days: **Acaramik (10ml/4gal) + Top Cop (15 ml/gal) + Foli-zyme (30ml/gal) + Stimulate (5ml/4 gal) + 10:55:10 HM (1Tbsp/gal)**

7-10 days: **Impetu (1Tbsp/gal) + Acaramik (10ml/4gal) + Sugar Mover (15ml/gal) + 5:10:27 HM (1Tbsp/gal)**

7-10 days: **Impetu (1Tbsp/gal) + Binder (5ml/gal) + Rezist (20ml/gal) + Bio-Forge (10ml/gal) + 5:10:27 HM (1Tbsp/gal)**

7-10 days: **Xstrata (5 ml/gal) + Top Cop (15ml/gal) + Binder (5ml/gal) + Forsa (2.5ml/gal) + Foli-zyme (30 ml/gal) + 5:5:45 HM (1Tbsp/gal)**

7-10 days: **Xstrata (5 ml/gal) + Ixus (5 ml/gal) + Sett (30ml/gal) + X-cyte (10ml/gal) + 5:10:27 HM (1Tbsp/gal)**

7-10 days: **Impetu (1Tbsp/gal) + Ixus (5 ml/gal) + Sett (30ml/gal) + Stimulate (5ml/4 gal) + 5:10:27 HM (1Tbsp/gal)**

MAFAS LTD

Lot 167, Bay View, Concord Road, La Romain. Trinidad W.I.
Tel: 868-354-9666; Tel/Fax: 868-697-1375. Email: marcusmycoo@yahoo.com
Visit us online: www.mafascaribbean.com

7-10 days: **Impetu** (1 TBSp/gal) + **Top Cop** (15ml/gal) + **Acaramik** (10ml/4gal) + **Sugar Mover** (20ml/gal) + **5:10:27 HM** (1Tbsp/gal)

7-10 days: **Diligent CO** (1Tbsp/gal) + **Acaramik** (10ml/4gal) + **Forsa** (5ml/gal) + **Flower Power** (20ml/gal) + **5:10:27 HM** (1Tbsp/gal)

7-10 days: **Diligent CO** (1Tbsp/gal) + **Binder** (5ml/gal) + **Sett** (30ml/gal) + **Bioforge** (10ml/gall) + **5:10:27 HM** (1Tbsp/gal)

7-10 days: **Xstrata** (5ml/gal) + **Binder** (5ml/gal) + **Foli-zyme** (30ml/gal) + **SugarMover (20 ml/ gal)** + **5:10:27 HM** (1Tbsp/gal)

7-10 days: **Xstrata** (5ml/gal) + **Top Cop** (15ml/gal) + **Acaramik** (5-10ml/4gal) + **Sugar Mover** (20ml/gal) + **5:5:45 HM** (1Tbsp/gal)

7-10 days: **Xstrata** (5ml/gal) + **Top Cop** (15ml/gal) + **Acaramik** (10ml/4gal) + **Forsa** (5 ml/gal) + **Flower Power** (20ml/gal) + **5:10:27 HM** (1Tbsp/gal)

7-10 days: **Diligent CO** (1Tbsp/gal) + **Binder** (5ml/gal) + **Sett** (30ml/gal) + **Bioforge** (10ml/gall) + **5:10:27 HM** (1Tbsp/gal)

7-10 days: **Diligent CO** (1TBSp/gal) + **Binder** (5ml/gal) + **Foli-zyme** (30ml/gal) + **5:10:27 HM** (1Tbsp/gal)

Weed Control: Best results used in Stoller pH+ treated water

- **Zibo 35 SC** (Glufosinate ammonium) for non-selective control of weeds. Best results when applied to young, actively growing weeds before they flower. Excellent for control of **White Top** @ 15 ml/gall
- **Icasso 24 EC** 2.5-5.0ml/gall for selective control of grasses including Fowl Foot Grass

Soil Applied fertilizer:

- Starter Fertiliser-**Pre-Plant** incorporate 2 bags **12-24-12** to soil or apply 1 1/2 oz/plant deep placed at planting
- Apply Soil Applied fertilizer 12:12:17+2 **every 4-6 weeks** or as needed
- Apply **Root Feed SP and Charge** as a **root drench every 14 days**. Dry 1 TBSp/plant or 1 TBSp/gall or 1-2 kg/50 gall barrel. Include 1-2 kg Harvest More for each stage of crop growth eg **10:55:10** (establishment), **5-10-27 Urea Mate** (flowering & fruiting)

Soil applied Fertilizers may need to be adjusted as the products increase efficiency of uptake. Review fertilizer application intervals and quantity.

MAFAS LTD

Lot 167, Bay View, Concord Road, La Romain. Trinidad W.I.
Tel: 868-354-9666; Tel/Fax: 868-697-1375. Email: marcusmycoo@yahoo.com
Visit us online: www.mafascaribbean.com

NOTES:

- Farmers are encouraged to contact **MAFAS LTD** personnel for further information on effective disease and pest control on crop production
- **X-tra Power, Stimulate, Root Feed SP** have strong NEMATICIDAL PROPERTIES when applied to the soil as a drench at the base of the plant at transplanting.
- **Top Cop** may be applied at 15ml/gal for control of a wide range of Bacterial and fungal leaf spots. Do a compatibility test. Can be mixed with **Sett, Nitro-plus 23, Stimulate** and **X-cyte** and other fungicides and Insecticides from Rotam.
- Always use **Stoller pH+** treated water in all sprays. Add **Ph+** (5ml/gal) first to your spray water .
- **Golden Pest Spray Oil** can be applied with all insecticide and fungicide mix as a pre-mix treatment. For control of eggs larval or nymphal stages of wide range pest
- **Golden Pest Spray Oil, Charge** and **Top Cop** are approved for use in ORGANIC CROPPING
- **Stimulate** is applied at knap sack rate of **5-10 ml/4 gall**
- Mist blower rates are twice the knapsack rate.
- **Sett, Nitro Plus, Root Feed** are essential to supply and maintain Calcium for fruit quality

Mite Control: Use **Acaramik** 10 ml/4 gall pre-mixed with **GPSO**@30 ml/gall then add water. Use at least twice at 5-7 days. If needed rotate with **Binder** 5ml/gall premixed with **GPSO** 30 ml/gall then add water

White Fly and Leafminer control: Apply **Rotaprid** @ 10 ml/gall as a drench 10-14 days after transplanting(see above)

During the crop: Foliar apply **Binder**@ 5ml/gall for -3 times. Rotate with **Rotaprid** @ 5ml/gall for 2-3 times

Gall Midge Control: Apply **Binder**@ 5ml/gall premixed with **GPSO** pre-mixed @ 30 ml/gall to flowers and fruit. Also drench below the plant (drip circle) with the same solution using a flat fan nozzle. Solution should be to at least a depth of 2 “in the soil.

Phytophthora Rot: Apply **Impetu** at 11/2TBSp/gall + **Bioforge** @ 10 ml/gall for every 5-7 days for three applications. Ensure full coverage. Rotate with any **Diligent** product for two applications. Always drench the soil if molded with **Aldabon**